

GENERATION AND ACCELERATION OF JETS FROM EFFECTIVE BOUNDARY LAYER AROUND BLACK HOLE

INDRANIL CHATTOPADHYAY AND SANDIP K. CHAKRABARTI

S.N. Bose National Centre for Basic Sciences, JD-Block, Salt Lake, Calcutta 700098

E-mail: indra@boson.bose.res.in and chakraba@boson.bose.res.in

We propose the processes of generation and radiative acceleration of jets from the centrifugal pressure dominated boundary layer around a black hole.

Proceedings of the 9th Marcel Grossman Meeting in Rome (Ed. R. Ruffini)

The inner boundary condition at the horizon for an inflow is that the radial flow velocity should be equal to the velocity of light. At an intermediate distance, centrifugal force becomes comparable to the gravity and matter is slowed down making it hot and puffed up like a tori. This is known as the centrifugal barrier supported boundary layer or CENBOL^{1,4}. This CENBOL acts as the effective boundary layer around black holes. Chakrabarti and Titarchuk¹ calculated the spectrum emitted from the CENBOL and showed that the CENBOL is the elusive ‘Compton cloud’ which gives rise to the power law hard tail. They also showed that if the accretion rate ($\dot{M}_d$) of the Keplerian component (the generator of soft photons) of the disc is greater than $\sim 0.4\dot{M}_{Edd}$, then the emitted spectrum will be without the hard tail due to thermal Comptonization and the black hole would be in the soft state. Subsequently it was realized^{2,3,4} that the CENBOL is not just a source of Comptonization but also acts as the source of jets and outflows. A part of the in falling matter is deflected by this effective boundary layer or CENBOL along the axis of symmetry to produce jets and outflows in at least some of the observed cases. This computed mass loss rates^{3,4} generally agree with the rates obtained through fully time dependent numerical simulations⁵. Recent reports on galactic black hole candidate GRS 1915+105⁶ and extragalactic black hole M87⁷ clearly show evidence that the region where jets originate and the region which emits Comptonized photons are the same. This vindicates our stand that the advective flow solution proposed more than ten years ago and which included accretion and winds interfaced by a CENBOL could be the correct paradigm.

Since CENBOL produces funnel wall as in a thick disk, and since matter and hard radiation can be thought of being emitted by the CENBOL surface, the effect of momentum deposited by the hot photons on outflows could be interesting to study. The effect of radiative momentum deposition on astrophysical outflows around compact objects were studied using early disc models by many workers (e.g. Vincent Icke⁸ and Abramowicz and Piran⁹). Chattopadhyay and Chakrabarti^{10,11}, revisited the problem using the radiation from advective discs¹. In this region, the subsonic outflowing matter is continuously bombarded by hot photons and hence apart from usual thermal acceleration, radiative acceleration should be important too. Thus the toroidal region or the CENBOL of accretion disc not only emits high energy radiation, but its typical shape helps the radiation to be focussed on the axis of symmetry. In our calculation of radiative momentum deposition force (RAMOD) we do not consider the attenuation of the radiation field due to interac-

tion with matter. With attenuation taken into account, optically thick flow would be difficult to accelerate this way. The two components of radiative momentum deposition force (RAMOD) have been calculated¹² and efforts are on include these forces in the analytical and numerical study. Our assumption of optically thin flow (i.e., $\tau < 1$) puts upper limit on the mass outflow rate and the mass of the central object^{11,12}:

$$\frac{\dot{M}^6}{M_B^5} < 1.7 \times 10^{-70} \text{ gm sec}^{-6}$$

or,

$$\dot{m}_{10}^6 m_{10} < 6.1 \times 10^{-8},$$

where, $\dot{m}_{10}$ is the mass outflow rate ($\dot{M}$) in the units of Eddington rate for a $10M_\odot$ black hole and m_{10} is the black hole mass (M_B) in units of $10M_\odot$. Increasing $\dot{M}_{acc}$ increases disk luminosity. This in turn increases the magnitude of RAMOD and the outflow rate $\dot{M}_{out}$. This increases the density ρ of the outflow thereby increasing τ and causing an attenuation of radiation by a factor of $e^{-\tau}$. Using the RAMOD we find the terminal speed of outflows achieved are about hundred percent higher than that achieved by purely thermally driven outflows. Of course, radiation drag force are seen to reduce the acceleration by some amount. But the drag is not very effective and we are unable to reach close to velocity of light by hydrodynamical processes alone. Thus our conclusion is that RAMOD does accelerate the outflow and higher RAMOD results in generally higher terminal speed. RAMOD also increases the energy of the flow often freeing bound matter (i.e., matter with negative energy) from the accretion disk.

This work is partly supported by DST project Analytical and Numerical Studies of Astrophysical Flows Around Compact Objects.

References

1. Chakrabarti S.K. and Titarchuk, L.G. *Astrophys. J.* **455**, 623 (1995)
2. S.K. Chakrabarti, *Ind. J. Phys.*, 72B, 183 (1998) astro-ph/9803227.
3. Chakrabarti, S. K. *Astron. Astrophys.*, **351** 185 (1999)
4. Das T.K. and Chakrabarti S.K. *Class. Quant. Grav.* **16**, 3879 (1999)
5. Molteni, D. Lanzafame, G. and Chakrabarti, S. K *Astrophys. J.*, **425** 161 (1994)
6. Fender, R. P. *astro-ph/9911176* (2000)
7. Junor, W. Biretta, J. and Livio, M. *Nature*, **401** 891 (1999)
8. Icke, V. *Astron. J.*, **85**(3) 329 (1980).
9. Abramowicz, M. A. and Piran, T. *it Astrophys. Journ.* **241** L7 (1980).
10. Chattopadhyay, I. and Chakrabarti, S. K. *Int. Journ. Mod. Phys. D*, **9**(1) 57 (2000).
11. Chattopadhyay, I. and Chakrabarti, S. K. *Int. Journ. Mod. Phys. D*, *in press*
12. Chattopadhyay, I. and Chakrabarti, S. K. in *Proc. of High Energy Gamma-Ray Astronomy* eds. F.A. Aharonian and H. Völk (in press)